

Math 314 Test 2, Example Problems #1

Michael Walton
Head Upper Division Tutor
BRIGHAM YOUNG UNIVERSITY MATH LAB

June 25, 2020

These questions are not from previous exams. These are questions that cover many, but not necessarily all, of the concepts which could appear on the exams. If you feel there is an error with the solutions, please contact the Math Lab via mathlabupper@mathematics.byu.edu, and we will rectify the mistake.

Selected Formulas

Cross Products: $a \times (b \times c) = (a \cdot c)b - (a \cdot b)c$

Second Derivative Test

Critical points: The point (a, b) is a critical point if $f_x(a, b) = 0$ and $f_y(a, b) = 0$ or they do not exist.

Given $D = f_{xx}(a, b)f_{yy}(a, b) - [f_{xy}(a, b)]^2$ and that (a, b) is a critical point, then:

- (a) If $D > 0$ and $f_{xx}(a, b) > 0$, then $f(a, b)$ is a local minimum.
- (b) If $D > 0$ and $f_{xx}(a, b) < 0$, then $f(a, b)$ is a local maximum.
- (c) If $D < 0$, then $f(a, b)$ is a saddle point.
- (d) If $D = 0$, we know nothing about $f(a, b)$.

True/False and Multiple Choice

1. There is a box with dimensions l, w, h . These dimensions change with time. At time $t = 7s$, the dimensions are $l = 3m$, $w = 2m$, $h = 6m$ and l and w are both increasing at a rate of $2m/s$ while h is decreasing at a rate of $5m/s$. Which of the following are decreasing at $t = 7s$?
 - A. Surface Area
 - B. Volume
 - C. Length of the diagonal
 - D. None of these
2. For the same box in the first question, which of these is the correct magnitude for the rate of change in surface area?
 - A. $9 \frac{m^2}{s}$
 - B. $18 \frac{m^2}{s}$
 - C. $36 \frac{m^2}{s}$
 - D. $59 \frac{m^2}{s}$
 - E. $118 \frac{m^2}{s}$
 - F. None of these

3. For the same box in the first question, which of these is the correct magnitude for the rate of change in volume?

A. $15 \frac{m^3}{s}$

B. $30 \frac{m^3}{s}$

C. $45 \frac{m^3}{s}$

D. $60 \frac{m^3}{s}$

E. $75 \frac{m^3}{s}$

F. $90 \frac{m^3}{s}$

G. None of these

4. For the same box in the first question, which of these is the correct magnitude for the rate of change in the length of the diagonal?

A. $-\frac{20}{7} \frac{m}{s}$

B. $-\frac{40}{7} \frac{m}{s}$

C. $-\frac{60}{7} \frac{m}{s}$

D. $-\frac{80}{7} \frac{m}{s}$

E. $-\frac{1000}{7} \frac{m}{s}$

F. None of these

5. Let $z = f(x, y)$ and both x and y are functions of t . Then

$$\frac{dz}{dt} = \frac{dz}{dx} \frac{dx}{dt} = \frac{dz}{dy} \frac{dy}{dt}$$

A. True B. False

6. If $6xy - \sin(x^2 + y) = x^2y^3$, what is $\frac{dy}{dx}$?

A. $\frac{dy}{dx} = \frac{2x \sin(x^2 + y) + 2xy^3 - 6y}{6x + 3x^2y^2 + \cos(x^2 + y)}$

B. $\frac{dy}{dx} = \frac{2x \sin(x^2 + y) + 2xy^3}{6x - 3x^2y^2 - \cos(x^2 + y)(2x + 1)}$

C. $\frac{dy}{dx} = \frac{2x \cos(x^2 + y) + 2xy^3}{6x + 3x^2y^2 + \cos(x^2 + y)}$

D. $\frac{dy}{dx} = \frac{2x \cos(x^2 + y) + 2xy^3 - 6y}{6x - 3x^2y^2 - \cos(x^2 + y)}$

E. $\frac{dy}{dx} = \frac{2x \sin(x^2 + y) + 2xy^3 - 6y}{6x - 3x^2y^2 - \cos(x^2 + y)(2x + 1)}$

F. None of these

7. What is the directional derivative of $f(x, y, z) = xe^y + ye^z + ze^x$ at the point $(0, 0, 0)$ in the direction $\langle 5, 1, -2 \rangle$?

A. $1/\sqrt{30}(5e^y + 5ze^x + e^z + xe^y - 2e^x - 2ye^z)$

B. $-1/\sqrt{30}(5e^y + 5ze^x + e^z + xe^y - 2e^x - 2ye^z)$

C. 0

D. $4/\sqrt{30}$

E. $-4/\sqrt{30}$

4. Find equations for the tangent planes for the following curves at the specified points.

(a) $z = x^2 \ln |x + y|$ at $(3, -2, 0)$

(b) $z = e^{x^2+y^2-1}$ at $(1, 0, 1)$

(c) $z = \cos((x^2 + y^2)\frac{\pi}{5})$ at $(4, -3, 1)$

5. Using the tangent planes found in the previous problem, estimate the z-coordinate of the following points of their respective curves.

(a) $z = x^2 \ln |x + y|$ at $(2.95, -2.1, z)$

(b) $z = e^{x^2+y^2-1}$ at $(1.1, -0.2, z)$

(c) $z = \cos((x^2 + y^2)\frac{\pi}{5})$ at $(4.2, -3.1, z)$

6. Evaluate $\int \int_R (5 - x) dA$, where $R = \{(x, y) : 0 \leq x \leq 5, 0 \leq y \leq 3\}$.

7. Find the absolute maximum and minimum values of $f(x, y) = xy^2$ on the set $D = \{(x, y) : x \geq 0, y \geq 0, x^2 + y^2 \leq 3\}$.

8. If $R = [-1, 3] \times [0, 2]$, use a Riemann sum with $m = 4$, $n = 2$ to estimate the value of $\int \int_R \sin(\pi(x + y)/6) dA$. Take the sample points to be the upper left corners of the squares.

9. Show that

$$0 \leq \int \int_R \sin(\pi x) \cos(\pi y) dA \leq \frac{1}{32}$$

where $R = [0, \frac{1}{4}] \times [\frac{1}{4}, \frac{1}{2}]$ by using the fact that

$$\int \int_D k dA = k(b-a)(d-c)$$

where $D = [a, b] \times [c, d]$ and k is a constant.

10. Calculate the following integral:

$$\int_0^1 \int_0^1 (u-v)^5 du dv$$

11. Find the mass and center of mass of an object that occupies the region $D = \{(x, y) : 0 \leq y \leq \sin(\pi x/L), 0 \leq x \leq L\}$ given that the density of the object is given by $\rho(x, y) = y$.

12. Find the average value of $f(x, y) = x^2y$ over the rectangle R whose vertices are $(-1,0)$, $(-1,5)$, $(1,5)$, $(1,0)$.

13. Use Lagrange multipliers to find the maximum and minimum values of the following functions:

(a) $f(x, y, z, t) = x + y + z + t$ under the constraint $x^2 + y^2 + z^2 + t^2 = 1$.

(b) $f(x, y, z) = \ln(x^2 + 1) + \ln(y^2 + 1) + \ln(z^2 + 1)$ under the constraint $x^2 + y^2 + z^2 = 12$.

(c) $f(x, y) = xe^y$ under the constraint $x^2 + y^2 = 2$.

14. Find the volume of the solid enclosed by the paraboloid $z = 2 + x^2 + (y - 2)^2$ and the planes $z = 1$, $x = 1$, $x = -1$, $y = 0$, and $y = 4$.